



DESCRIPTION

A broad-spectrum nematicide and taenicide anthelmintic, endoparasiticide.

INDICATIONS

- Fenbendazole has a broad-spectrum of activity against gastrointestinal roundworms and lungworms of livestock, including adults and L4-larvae of the most important species (e.g. of the genus *Bunostomum*, *Haemonchus*, *Ostertagia*, *Teladorsagia*, *Trichostrongylus*, *Cooperia*, *Nematodirus*, *Chabertia*, *Oesophagostomum*, *Trichuris*, *Dictyocaulus*, *Muellerius*, etc.) as well as arrested larvae of several species.
- It is also effective against most livestock tapeworms (e.g. *Moniezia*, *Taenia*).
- In horses it controls the major parasitic roundworms such as Large Strongyles (*Cyathostomins*), Small Strongyles (*Strongylus* spp), *Parascaris equorum*, etc. as well as tapeworms. (e.g. *Anaplocephala* spp).
- It is also effective against the major parasitic roundworms (e.g. *Ancylostoma*, *Toxocara*, *Trichuris*, *Uncinaria*) and tapeworms (e.g. *Echinococcus*, *Dipylidium*, *Taenia*, etc.) of dogs and cats.

CONTENTS:

Each ml contain
Fenbendazole I.P (Vet) 25 mg.

Dose:

Dog & Cat : 25 - 50 mg/kg body weight
Goat & sheep : 5-15 mg/kg for few parasites upto 125-150 mg/kg body weight
Pig : 2-25 mg/kg body weight
Cattle : 5-15 mg/kg body weight
Horse : 5-15 mg/kg for few parasites upto 150 mg/kg body weight
Poultry : 10-50 mg/kg body weight

Or as recommended by the Veterinarian