Nuvectin



DESCRIPTION

Ivermectin is a macrocyclic lactone anthelmintic used to treat a variety of internal and external parasites commonly found in animals and birds.

INDICATIONS

Poultry: Most species of adult roundworms, including Ascaridia galli (large roundworm), Heterakis gallinarum (cecal worm). Syngamus trachea (gapeworm), and Oxyspirura mansoni (manson eyeworm). It has variable effectiveness against Capillaria (threadworms) at the dosage that can be safely administered. Some species of mites, including Cytodites nudus (airsac mites). Laminosioptes cisticola (fowl cyst mites). Knemidocoptes mutans (scaly leg mites), and Micnemidocoptes derool. It has mixed effectiveness with management of Ornithonyssus spp (fowi mites).

Cattle & Swine: Gastrointestinal roundworms (adults and 4th stage larva), lungworms (adult and 4th stage larva), cattle grubs (parasitic stages), sucking lice, eye worms (Thelazia) and miles (scabies).

Camel: Gastrointestinal nematodes, scabies and filariasis.

Sheep: Gastrointestinal roundworms (adults and 4th stage larva), lungworms (adults and 4th stage larva), nasal bots and mange mites (sheep scab).

CONTENTS:

Each ml contain Ivermectin BP 0.8 mg Benzyl alcohol BP 2 % as preservative

Dose:

Poultry: 0.2 mg/kg Body weight or 350 - 400 ml / 1000 birds (layers).

Cattle, Horses, Swine, Sheep & Goat:

0.2 mg/kg body weight or 10 ml/40 kgs Body weight.

Dogs & Cats: 0.2 mg/kg Body weight

Nuvectin



Rabbits: 0.2 mg to 0.3 mg/kg body weight

OR AS RECOMMENDED BY THE VETERINARIAN